

Food Attractants to Increase Pheromone-Baited Trap Performance for *Scyphophorus acupunctatus* (Coleoptera: Dryophthoridae) in Mezcal Maguey

Author(s): Pedro Figueroa-Castro, Víctor López-Martínez, Francisca Silva-García and Héctor González-Hernández

Source: Florida Entomologist, 100(1):203-205.

Published By: Florida Entomological Society

DOI: <http://dx.doi.org/10.1653/024.100.0135>

URL: <http://www.bioone.org/doi/full/10.1653/024.100.0135>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Food attractants to increase pheromone-baited trap performance for *Scyphophorus acupunctatus* (Coleoptera: Dryophthoridae) in mezcal maguey

Pedro Figueroa-Castro¹, Víctor López-Martínez^{1,*}, Francisca Silva-García², and Héctor González-Hernández³

Agave weevil, *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Dryophthoridae), is the main pest on wild and cultivated agave in Mexico (González-Hernández et al. 2007; Aquino Bolaños et al. 2014). In Guerrero State, this weevil infests wild, semi-cultivated, and cultivated maguey for producing mezcal (*Agave vivipara* L. = *Agave angustifolia* Haworth, and *Agave cupreata* Trel. & Berger; Asparagaceae) (Barrios Ayala et al. 2006; P. F.-C. unpublished data). Conventional management tactics (i.e., pesticide applications) have been inefficient to control agave weevil populations mainly because of the cryptic habits of this weevil, which completes its life cycle inside the agave boll (P. F.-C. unpublished data).

A pheromone-baited trap system is the standard tool for monitoring adult populations of *S. acupunctatus* on blue agave in Mexico (Figueroa-Castro et al. 2013). The use of agave tissue as a bait was proposed by Rojas et al. (2006). Further, Figueroa-Castro (2014) confirmed the need to bait pheromone traps with agave tissue (300–400 g) to synergize the response of agave weevils to pheromone-baited traps, increasing the number of captured weevils.

However, this system has some inconvenience. Firstly, although agave tissues may be obtained directly from infested agave plants, the recommended practices to eliminate infested plants can make it difficult to find them, and in this case it is necessary to obtain the tissues from healthy agave plants with a commercial value (plants in growth stages, shoots from rhizomes of parental plants, termed “hijuelos”). Secondly, time is required for plant selection, dissection, and for preparing the bait. Thirdly, each plant with a commercial value but used as bait represents a future loss in mezcal yield. With the objective to find an efficient, economic, and practical food bait, 2 alternative food attractants (banana and pineapple ripe fruits) were evaluated to synergize the response of agave weevil to synthetic pheromone-baited traps in the mezcal maguey area.

Traps of the type TOCCI (4 L white buckets) used in this study were handmade (Figueroa et al. 2016a). Each trap was baited with a dispenser of synthetic aggregation pheromone of *S. acupunctatus* (Tequilur[®], Ferro-CompS, Mexico City, Mexico) and 300 g of the food attractant. The food attractant was placed inside a 0.5 L plastic container with holes (0.5 cm diameter) and sprinkled with malathion insecticide (Malathion 1000[®], Anajalsa, S. A. de C. V., Jalisco, Mexico) (10 mL/L of water). The pheromone dispenser was changed monthly and the food attractant every 15 d.

The experiment was performed from 29 Jan to 21 May 2016 in a 4-yr-old espadín maguey (*A. vivipara*, *A. angustifolia*) plantation, in El Rancho Frío (18.38678°N, 99.16611°W; 1,120 m asl), located in Quetzalapa, Huitzuc de los Figueroa, Guerrero, Mexico. We evaluated the following treatments: 1) empty trap without synthetic pheromone and food attractant but with soapy water (control), 2) trap baited with pheromone but without food attractant, 3) trap baited with pheromone and 300 g of vegetative tissue of papalote maguey, 4) trap baited with pheromone and 300 g of vegetative tissue of espadín maguey, 5) trap baited with pheromone and 300 g of ripe banana, and 6) trap baited with pheromone and 300 g of ripe pineapple.

Traps were placed at ground level, near an agave plant, spatially distributed in a square pattern (as recommended by Figueroa-Castro et al. 2016b), with a distance between traps of 50 m. The experiment was established in a completely randomized experimental design, each treatment with 4 replications. Traps were checked biweekly for 5 sample dates. On each visitation date, we emptied the traps and collected the captured weevils. Specimens were taken to the laboratory of Agrícola Production at the Universidad Autónoma del Estado de Morelos (Cuernavaca, Morelos, Mexico) to be counted and sexed based on the morphology of the final abdominal segment (Ramírez-Choza 1993).

Statistical analysis was performed using SAS/STAT[®] software (SAS Institute 2004). The data were analyzed by analysis of variance (ANOVA) using proc GLM. Prior to ANOVA, data were checked for normality (Shapiro–Wilk test) and homogeneity of variances (Bartlett test), and data transformation was not necessary. Treatment means were separated with a Tukey test ($\alpha = 0.05$). A chi-squared test was applied to determine differences in the number of females and males caught in traps.

The number of agave weevils captured was affected by the treatments, and we found statistical differences between treatments ($F = 14.30$; $df = 5,15$; $P < 0.0001$). Traps baited with the synthetic pheromone and any food bait (papalote maguey tissue, banana, or pineapple) captured significantly more weevils than traps baited only with pheromone. Empty traps did not capture weevils (Fig. 1).

The fact that host plant volatiles synergize the responses of weevils to the synthetic pheromone has been reported for several species (Tinzaara et al. 2007; Wibe et al. 2014). With regard to food attractants, the best food attractants in our study were papalote maguey tissue and

¹Universidad Autónoma del Estado de Morelos, Facultad de Ciencias Agropecuarias, Cuernavaca, Morelos, CP 62209, México, E-mail: figueroac.pedro@gmail.com (P. F.-C.), victor.lopez@uaem.mx (V. L.-M.)

²Universidad Autónoma Chapingo, Posgrado en Desarrollo Rural Regional, Chapingo, Estado de México, CP 56230, México; E-mail: fran_silva19@yahoo.com.mx (F. S.-G.)

³Colegio de Postgraduados, Postgrado en Fitosanidad-Entomología y Acarología, Montecillo, Texcoco, Edo. de México, CP 56230, México; E-mail: hgzzhdz@colpos.mx (H. G.-H.)

*Corresponding author; E-mail: victor.lopez@uaem.mx (V. L.-M.)




Fig. 1. Mean number of *Scyphophorus acupunctatus* weevils captured per trap biweekly, with various food attractants. Means were calculated from data of 5 biweekly samples. Treatments with similar letters are not significantly different (Tukey, $\alpha = 0.05$). Error bars indicate SE.

ripe banana and pineapple fruit; these fruit baits have been reported previously as food attractants for trapping other weevil species. Pineapple was evaluated by Valdés et al. (2005) and García-Ramírez et al. (2014) for trapping *S. acupunctatus* on tuberose plant (*Polianthes tuberosa* L. 'Perla' [Asparagaceae]). Figueroa-Castro et al. (2016b,c) used agave tissue for collecting *S. acupunctatus* in mezcal maguey and in blue agave. Bananas were tested by Al-Saoud (2011) for collecting *Rhynchophorus ferrugineus* (Olivier), and by Sumano et al. (2012) for trapping *Rhynchophorus palmarum* (L.) (Coleoptera: Dryophthoridae). Banana and pineapple fruit appear to be good food baits for the agave weevil, with similar captures as those reported for agave tissues, but with additional advantages. They 1) are cheaper than agave, 2) are abundant, 3) are easy to buy, and 4) have no direct impacts on agave production.

Significantly more *S. acupunctatus* females than males were captured in all the treatments. These findings were similar to previous reports (Macedonio-Guevara 2015; Figueroa-Castro et al. 2016c). Capturing a large number of females is very important, because it has a direct impact for decreasing future population densities of agave weevil.

Based on our results, for monitoring agave weevil populations on any cultivated agave we suggest baiting the traps with pheromone and 300 g of food attractants such as papalote maguey tissue, ripe banana, or ripe pineapple fruit. Future research should focus on the potential to use this trapping system for initiating management practices, for preventing establishment of agave weevil populations in new plantations, and as part of a mass trapping system for controlling the population density of this agave pest.

We thank Daniel Castro Rodríguez (mezcal maguey grower) for access to all the facilities on the maguey plantation in Quetzalapa, Guerrero. We also thank Luis Figueroa Ocampo and Daniel Castro Rodríguez for their valuable assistance in the establishment and evaluation of field experiments. In addition, we thank the Programa para el Desarrollo Profesional Docente (SEP PRODEP) for providing a postdoctoral scholarship to the first author at the Facultad de Ciencias Agropecuarias, Universidad Autónoma del Estado de Morelos. The third author acknowledges the Consejo Nacional de Ciencia y Tecnología (CONACYT) scholarship for M.Sc. graduate studies.

Summary

The agave weevil, *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Dryophthoridae), is the main insect pest on maguey grown for mezcal beverage production in Guerrero, Mexico. We studied the effectiveness of several food attractants in pheromone-baited traps for

capturing *S. acupunctatus* in mezcal maguey plantations. All of the tested food attractants synergized the response of the agave weevil to synthetic aggregation pheromone traps. Adding papalote maguey tissue, ripe banana, or ripe pineapple fruit increased the number of captured weevils in pheromone-baited traps. However, it is easier, cheaper, and more practical to use ripe banana or pineapple fruit as the bait attractant than using the agave tissue.

Key Words: agave weevil; food attractant; monitoring; pheromone trap

Sumario

El picudo del agave *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Dryophthoridae), es la principal plaga en maguey mezcalero en Guerrero, México. En la presente investigación se evaluó la efectividad de varios atrayentes alimenticios en trampas con feromona de agregación sintética para la captura de *S. acupunctatus* en plantaciones de maguey mezcalero espadín. Todos los atrayentes alimenticios empleados potencializan la respuesta de este picudo a las trampas con feromona sintética. Colocar tejidos de maguey papalote, fruta madura de plátano o piña incrementa el número de picudos capturados en las trampas con feromona. Sin embargo, emplear fruta madura de plátano o piña es más sencillo, económico y práctico que usar tejido de agave.

Palabras Clave: picudo del agave; atrayentes alimenticios; monitoreo; trampas con feromona

References Cited

- Al-Saoud AH. 2011. Comparative effectiveness of four food baits in aggregation pheromone traps on red palm weevil *Rhynchophorus ferrugineus* Olivier. Arab Journal of Plant Protection 29: 83–89.
- Aquino Bolaños T, Pozo Velázquez E, Álvarez Hernández U, Delgado Gamboa JR. 2014. Plantas hospedantes del picudo del agave *Scyphophorus acupunctatus* (Gyllenhal) (Coleoptera: Curculionidae) en Oaxaca, México. Southwestern Entomologist 39: 163–169.
- Barrios Ayala A, Ariza Flores R, Molina Muñoz JM, Espinosa Paz H, Bravo Mosqueda E. 2006. Manejo de la fertilización en magueyes mezcaleros cultivados (*Agave* spp.) de Guerrero. Folleto Técnico No.13. Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias, Iguale, Guerrero, Mexico.
- Figueroa-Castro P. 2014. Factores asociados a la trampa y al atrayente, que influyen en la eficiencia del sistema de trapeo para el monitoreo del picudo del agave. Unpublished doctoral thesis. Postgrado en Fitosanidad, Colegio de Postgraduados, Montecillo, Mexico.
- Figueroa-Castro P, Solís-Aguilar JF, González-Hernández H, Rubio-Cortés R, Herrera-Navarro EG, Castillo-Márquez LE, Rojas JC. 2013. Population dynamics of *Scyphophorus acupunctatus* (Coleoptera: Curculionidae) on blue agave. Florida Entomologist 96: 1454–1462.
- Figueroa-Castro P, López-Martínez V, Hernández-Ruiz A, Silva-García F, Campos-Figueroa M. 2016a. Determining the best pheromone-baited traps for capturing *Scyphophorus acupunctatus* (Coleoptera: Dryophthoridae) in mezcal agave. Florida Entomologist 99: 790–792.
- Figueroa-Castro P, González-Hernández H, Carrillo-Sánchez JL, Solís-Aguilar JF, del Real-Laborde JI, Rojas JC. 2016b. Effect of the height and distribution pattern of pheromone-baited traps on the capture of *Scyphophorus acupunctatus* (Coleoptera: Dryophthoridae) on blue agave. Florida Entomologist 99: 297–299.
- Figueroa-Castro P, Rodríguez-Rebollar H, González-Hernández H, Solís-Aguilar JF, del Real-Laborde JI, Carrillo-Sánchez JL, Rojas JC. 2016c. Attraction range and inter-trap distance of pheromone-baited traps for monitoring *Scyphophorus acupunctatus* (Coleoptera: Dryophthoridae) on blue agave. Florida Entomologist 99: 94–99.
- García-Ramírez MJ, López-Martínez V, Alia-Tejagal I, Andrade-Rodríguez M, Rojas JC. 2014. Influence of trap color and food bait on the catches of *Scyphophorus acupunctatus* by pheromone-baited traps in tuberose crop. Journal of the Kansas Entomological Society 87: 96–101.
- González-Hernández H, Solís Aguilar JF, Pacheco-Sánchez C, Flores-Mendoza FJ, Rubio-Cortés R, Rojas JC. 2007. Insectos barrenadores del agave tequilero, pp. 39–67 In González-Hernández H, del Real-Laborde JI, Solís Aguilar

- JF [eds.], Manejo de Plagas del Agave Tequilero. Colegio de Postgraduados and Tequila Sauza S.A. de C.V., Zapopan, Jalisco, México.
- Macedonio-Guevara A. 2015. Abundancia estacional del picudo del agave en plantaciones de agave mezcalero en Guerrero. Unpublished M.Sc. thesis. Postgrado en Fitosanidad, Colegio de Postgraduados, Montecillo, Mexico.
- Ramírez-Choza JL. 1993. Max del henequén *Scyphophorus interstitialis* Gylh. Bioecología y control. Serie: Libro Técnico. Centro de Investigación Regional del Sureste. Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias, Secretaría de Agricultura, Ganadería y Recursos Hidráulicos, Mérida, Yucatan, Mexico.
- Rojas J, González-Hernández H, Ruiz-Montiel C, Rangel-Reyes DN, Ceja EI, García-Coapio G, del Real-Laborde I. 2006. Optimización de un sistema de monitoreo/trampeo masivo para el manejo del picudo del agave, *Scyphophorus acupunctatus* Gylh, pp. 51–58 In Barrera JF, Montoya P [eds.], Simposio sobre trampas y atrayentes en detección, monitoreo y control de plagas de importancia económica. Sociedad Mexicana de Entomología/El Colegio de la Frontera Sur. Manzanillo, Colima, Mexico.
- SAS Institute Inc. 2004. SAS/STAT® 9.1 User's Guide. SAS Institute Inc., Cary, North Carolina.
- Sumano D, Sánchez S, Romero J, Sol A. 2012. Eficacia de captura de *Rhynchophorus palmarum* L. (Coleoptera: Dryophthoridae) con diferentes diseños de trampas en Tabasco, México. Fitosanidad 16: 43–78.
- Tinzaara W, Gold CS, Dicke M, Huis AV, Ragama PE. 2007. Host plant odours enhance the responses of adult banana weevil to the synthetic aggregation pheromone Cosmolure+®. International Journal of Pest Management 53: 127–137.
- Valdés EM, Lucila Aldana L, Figueroa BR, Gutiérrez OM, Hernández RMC, Tomasa M. 2005. Trapping of *Scyphophorus acupunctatus* (Coleoptera: Curculionidae) with two natural baits in a field of *Polianthes tuberosa* (Liliales: Agavaceae) in the state of Morelos, Mexico. Florida Entomologist 88: 338–340.
- Wibe A, Borg-Karlson AK, Cross J, Bichao H, Fountain M, Liblikas I, Sigsgaard L. 2014. Combining 1,4-dimethoxybenzene, the major flower volatile of wild strawberry *Fragaria vesca*, with the aggregation pheromone of the strawberry blossom weevil *Anthonomus rubi* improves attraction. Crop Protection 64: 122–128.